The table below includes weighted estimates of variant proportions by state/jurisdiction collected over the **four weeks ending 1/22/2022**.
This analysis represents a different collection period and length of time, four weeks, compared to the weighted estimates shown above at the national and regional level which are calculated by week.

State/Jurisdiction	Total Sequences	B.1.1.529 (Omicron)	BA.2 (Omicron)	B.1.617.2 (Delta)	AY.1 (Delta)	Other
Alabama	937	97.35% (96.1-98.3%)	0.12%† (0.0-0.6%)	1.99% (1.2-3.1%)	0.00%† (NA)	0.54%† (0.2-1.3%)
Alaska	701	94.02% (92.0-95.7%)	1.21%† (0.2-3.7%)	4.50% (2.2-8.0%)	0.00%† (NA)	0.26%† (0.0-2.2%)
American Samoa	0					
Arizona	3,330	93.37% (91.9-94.6%)	0.42%† (0.0-2.0%)	6.11% (4.4-8.2%)	0.00%† (NA)	0.10%† (0.0-0.7%)
Arkansas	1,044	95.18% (91.3-97.7%)	0.10%† (0.0-0.8%)	4.60% (2.3-8.2%)	0.00%† (NA)	0.13%† (0.0-1.1%)
California	23,442	96.80% (95.1-98.0%)	0.23% (0.2-0.3%)	2.88% (1.8-4.5%)	0.00%† (NA)	0.09%† (0.0-0.4%)
Colorado	5,394	97.54% (95.9-98.6%)	0.11%† (0.0-0.2%)	2.08% (1.1-3.5%)	0.00%† (NA)	0.27%† (0.0-1.5%)
Commonwealth of the Northern Mariana Islands	76	50.00%† (NA)	1.32%† (NA)	48.69%† (NA)	0.00%† (NA)	0.00%† (NA)
Connecticut	2,215	95.13% (90.7-97.8%)	0.06%† (0.0-0.3%)	4.53% (2.3-8.0%)	0.00%† (NA)	0.27%† (0.0-2.1%)
Delaware	1,507	95.78% (94.6-96.7%)	0.06%† (0.0-0.7%)	4.03% (3.1-5.2%)	0.00%† (NA)	0.13%† (0.0-1.6%)
District of Columbia	654	98.70% (97.5-99.4%)	0.20%† (0.0-0.9%)	1.10%† (0.4-2.2%)	0.00%† (NA)	0.00%† (NA)
Federated States of Micronesia	0					
Florida	7,749	98.16% (97.0-99.0%)	0.05%† (0.0-0.2%)	1.28% (1.0-1.6%)	0.00%† (NA)	0.51%† (0.0-2.3%)
Georgia	5,706	98.26% (96.7-99.2%)	0.02%† (0.0-0.1%)	1.50% (0.8-2.6%)	0.00%† (NA)	0.21%† (0.0-1.2%)
Guam	72	99.23%† (65.4-100.0%)	0.00%† (NA)	0.77%† (0.0-34.6%)	0.00%† (NA)	0.00%† (NA)
Hawaii	878	99.14% (98.3-99.6%)	0.39%† (0.1-1.4%)	0.42%† (0.1-1.1%)	0.00%† (NA)	0.05%† (0.0-0.5%)
Idaho	2,819	93.81% (91.6-95.6%)	0.07%† (0.0-0.3%)	6.04% (4.3-8.3%)	0.00%† (NA)	0.07%† (0.0-0.6%)
Illinois	3,976	92.68% (88.2-95.8%)	0.03%† (0.0-0.2%)	6.55% (3.9-10.3%)	0.02%† (0.0-0.1%)	0.72%† (0.0-3.2%)
Indiana	1,802	89.91% (82.7-94.8%)	0.06%† (0.0-0.4%)	9.80% (5.2-16.4%)	0.00%† (NA)	0.23%† (0.0-1.4%)
Iowa	1,547	86.07% (76.0-93.0%)	0.00%† (NA)	13.66% (6.8-23.7%)	0.05%† (0.0-0.3%)	0.21%† (0.0-1.1%)
Kansas	1,685	88.24%	0.06%†	11.51%	0.00%†	0.19%†

Table includes weighted estimates of variant proportions from sequences collected over the **four weeks ending 1/22/2022** obtained through CDC genomic surveillance (NS3 and CDC-funded sequencing contracts) and tagged baseline surveillance sequencing submitted to public repositories by state, local, academic, or commercial laboratories. The four week timeframe used for the analysis will roll forward each week. Variants with zero observations in all jurisdictions are not enumerated in this chart (B.1.1.7/Alpha, B.1.351/Beta, P.1/Gamma, P.2/Zeta, B.1.427/429/Epsilon, B.1.525/Eta, B.1.526/Iota, B.1.617.1/Kappa, B.1.617.3, AY.2/Delta, B.1.621/Mu).
* Other represents weighted estimates for variants not designated as a variant of concern (VOC), variant of interest (VOI), or variant being monitored (VBM).
† Estimates are less reliable based on one or more violations of NCHS data presentation standards for proportions: number of sequences submitted by state in four-week time period was ..

Download Data

Updated February 15, 2022