The table below includes weighted estimates of variant proportions by state/jurisdiction collected over the **four weeks ending 3/26/2022**.
This analysis represents a different collection period and length of time, four weeks, compared to the weighted estimates shown above at the national and regional level which are calculated by week.

State/Jurisdiction	Total Sequences	B.1.1.529 (Omicron)	BA.2 (Omicron)	BA.2.12.1 (Omicron)	B.1.617.2 (Delta)	B.1.1.7 (Alpha)	Other
Alabama	69	74.09% (62.1-83.9%)	25.91% (16.1-37.9%)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
Alaska	386	37.15%† (21.0-55.7%)	62.85%† (44.3-79.0%)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
American Samoa	0						
Arizona	979	73.99% (69.5-78.2%)	25.51% (21.4-30.0%)	0.38%† (0.0-1.4%)	0.11%† (0.0-0.6%)	0.00%† (NA)	0
Arkansas	41	80.64%† (60.0-93.6%)	19.36%† (6.4-40.0%)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
California	6,842	65.67% (62.2-69.0%)	34.12% (30.9-37.5%)	0.17%† (0.0-0.5%)	0.04%† (0.0-0.1%)	0.00%† (NA)	0
Colorado	1,861	70.00% (63.5-75.9%)	28.93% (23.1-35.4%)	0.93% (0.5-1.5%)	0.14%† (0.0-1.5%)	0.00%† (NA)	0
Commonwealth of the Northern Mariana Islands	162	9.88%† (NA)	90.12%† (NA)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
Connecticut	919	51.17%† (34.9-67.3%)	46.59%† (30.0-63.7%)	2.24%† (0.2-9.2%)	0.00%† (NA)	0.00%† (NA)	0
Delaware	146	69.04%† (49.7-84.5%)	30.96%† (15.5-50.3%)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
District of Columbia	297	44.82% (36.0-53.9%)	53.43% (46.1-60.7%)	1.75%† (0.2-5.9%)	0.00%† (NA)	0.00%† (NA)	0
Federated States of Micronesia	0						
Florida	3,445	58.82% (56.6-61.0%)	40.42% (38.2-42.7%)	0.76% (0.5-1.1%)	0.00%† (NA)	0.00%† (NA)	0
Georgia	859	68.71% (60.0-76.6%)	30.38% (22.5-39.2%)	0.69%† (0.0-3.7%)	0.22%† (0.0-1.0%)	0.00%† (NA)	0
Guam	11	0.00%† (NA)	100.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
Hawaii	518	43.89% (33.1-55.1%)	55.91% (44.6-66.8%)	0.20%† (0.0-1.1%)	0.00%† (NA)	0.00%† (NA)	0
Idaho	184	83.56% (76.8-89.0%)	15.75% (10.8-21.8%)	0.69%† (0.0-8.7%)	0.00%† (NA)	0.00%† (NA)	0
Illinois	2,214	56.72% (51.2-62.1%)	41.81% (36.8-46.9%)	1.00% (0.6-1.6%)	0.34%† (0.0-1.7%)	0.00%† (NA)	0 (0.0
Indiana	405	72.15% (66.9-77.0%)	27.08% (21.9-32.8%)	0.54%† (0.0-4.1%)	0.23%† (0.0-2.3%)	0.00%† (NA)	0
Iowa	344	84.75% (75.7-91.4%)	14.62% (8.7-22.5%)	0.00%† (NA)	0.63%† (0.1-2.2%)	0.00%† (NA)	0
Kansas	132	76.86% (67.0-85.0%)	23.14% (15.0-33.0%)	0.00%† (NA)	0.00%† (NA)	0.00%† (NA)	0
Kentucky	368						

Table includes weighted estimates of variant proportions from sequences collected over the **four weeks ending 3/26/2022** obtained through CDC genomic surveillance (NS3 and CDC-funded sequencing contracts) and tagged baseline surveillance sequencing submitted to public repositories by state, local, academic, or commercial laboratories. The four week timeframe used for the analysis will roll forward each week. Variants with zero observations in all jurisdictions are not enumerated in this chart (B.1.351/Beta, P.1/Gamma, P.2/Zeta, B.1.427/429/Epsilon, B.1.525/Eta, B.1.526/Iota, B.1.617.1/Kappa, B.1.617.3, AY.1/Delta, AY.2/Delta, B.1.621/Mu). Pango lineages Q.1, Q.2, and Q.3 are sublineages of Alpha B.1.1.7 and are included in the proportions of B.1.1.7. Delta B.1.617.2 sublineages AY.3 - AY.133 and their sublineages are included in the parental B.1.617.2. Omicron sublineages BA.1, BA.3, BA.4, BA.5 and their sublineages are aggregated with B.1.1.529. Except BA.2.12.1, BA.2 sublineages are aggreagated..

Download Data

Updated April 19, 2022